

USPAS Graduate Accelerator Physics Homework 6

Due date: Wednesday January 30, 2019

1 LEP electron radiation

Consider 50 GeV electrons in LEP (*requiesce in pace*), circulating in a nearly-circular ring with a total circumference of about 27 km. Consider also a 5 TeV electron storage ring that is built around the earth's equator. For each accelerator:

- How much energy is radiated per electron per turn?
- What is the characteristic damping time τ_0 , in turns and in seconds?
- What is the bending field?

2 RHIC gold ion radiation

In RHIC, 55 bunches of 10^9 gold ions with $(Z, A) = (79, 197)$ circulate in each ring, at a top energy of $\gamma = 108$. The circumference is 3833 m and the main arc dipole bending radius is $\rho = 243$ m.

- How much energy is radiated per gold ion per turn?
- What is the characteristic damping time τ_0 , in turns and in seconds?
- The synchrotron radiation power is a serious cryogenic load if it exceeds about 1 W/m. Is it a problem?
- If 360 bunches of 2×10^{11} 10 GeV electrons circulate in a new ring in the existing RHIC tunnel, how much energy is radiated per electron per turn, and what is the characteristic damping time τ_0 ?

3 FCC wall-plug power

A Future Circular Collider (FCC) might circulate 50 TeV protons in a 100 km circumference tunnel with a main arc dipole bend radius of 11 km.

- What is the arc dipole bending field?
- What is the critical energy of photons radiated in the dipoles?
- What is the total energy lost per turn, per proton?
- If each proton beam has a current of 0.5 A, what is the total synchrotron radiation power, per ring?
- Assuming that cryogenic refrigerators operate with a Carnot efficiency of 20%, how much "wall-plug" power would be required if the synchrotron radiation were absorbed at a temperature of 4 K?

4 Muon storage ring

Consider muon storage rings.

- (a) What is the total power radiated in an isomagnetic muon storage ring of radius ρ with an average beam current of I ?
- (b) How much power is emitted in a 30 GeV ring with $\rho = 250$ m that stores 1 A of electrons, or 1 A of muons?
- (c) What is the natural beam current lifetime decay for 30 GeV muons?
- (d) What is the heat load due to the decay of 1 A of 30 GeV muons?

5 Light source calculations

A light source with a circumference of 176 m contains 8 identical DBA cells, with zero dispersion (and dispersion slope) at one end or the other of all 16 dipoles. Each dipole is 2.7 m long, and the beam energy is 2.5 GeV.

- (a) What is the characteristic energy of the photons radiated in the dipoles?
- (b) How much energy is radiated per turn, per electron?
- (c) What is the momentum compaction factor of the ring?
- (d) What are the damping times τ_x , τ_y , and τ_s ?
- (e) What is the natural horizontal emittance (approximately)?