

# USPAS Accelerator Physics 2019

## Northern Illinois University and UT-Batelle

### Octupoles, Detuning, and Slow Extraction

Todd Satogata (Jefferson Lab and ODU) / [satogata@jlab.org](mailto:satogata@jlab.org)



Steve Peggs (BNL) / [peggs@bnl.gov](mailto:peggs@bnl.gov)

Bhawin Dhital (ODU) / [bdhit001@odu.edu](mailto:bdhit001@odu.edu) and Kiel Hock (BNL) / [khock@bnl.gov](mailto:khock@bnl.gov)

<http://www.toddsatogata.net/2019-USPAS>

Happy Birthday to Elijah Wood, Sarah McLachlan, Alan Alda, and Jackson Pollack!  
Happy National Blueberry Pancake Day, Data Privacy Day, and National Kazoo Day!

# Spring Break Adventures: McKay's



<http://www.mckaybooks.com>





## Overview (Afternoon)

- Useful nonlinearities
- Octupoles and detuning
- Discrete motion in  $(J, \phi)$  space
  - Difference Hamiltonian
  - More lecturer self-indulgence
- Motion near half-integer tunes
  - Contours of constant Hamiltonian (energy)
- Half-integer slow extraction
  - A useful application of first-order octupole perturbation theory
- Extending to third-integer extraction
- Modern use: resonance island extraction at CERN



# Useful Nonlinearities

- Catch-22 revisited
  - Nonlinearities are unavoidable in accelerators
  - Nonlinearities can correct motion – to a degree
  - Nonlinearities add higher “order” nonlinear behavior
  - But nonlinearities can be used for good!
  - Octupoles introduce new first-order behavior



## Integrable Particle Dynamics in Accelerators

### Sponsors:

Northern Illinois University and UT-Battelle

### Course Name:

Integrable Particle Dynamics in Accelerators

### Instructor:

Sergei Nagaitsev and Timofey Zolkin, Fermilab

This  
USPAS  
session!




# 1D Single Octupole Kick

$(x_p, x'_p)$  : physical coordinates  
 $(x, x')$  : normalized coordinates

$$\begin{pmatrix} x_p \\ x'_p \end{pmatrix} = \begin{pmatrix} \sqrt{\beta} & 0 \\ -\alpha/\sqrt{\beta} & 1/\sqrt{\beta} \end{pmatrix} \begin{pmatrix} x \\ x' \end{pmatrix}$$


inverse Floquet transformation, book Equation 3.22

$$M = I \cos(2\pi Q_0) + J \sin(2\pi Q_0)$$



$$\Delta x'_p = -g_p x_p^3$$

$$g_p \equiv \frac{B''' L}{B\rho} \text{ (be careful)}$$



- Linear 1D lattice with single octupole kick

$$\Delta x' = -g x^3 \quad g \equiv g_p \beta^2$$

# 1D Single Octupole Detuning and Resonances

$$\Delta x' = -gx^3 \quad g \equiv g_p \beta^2$$

- Use the normalized phase space figure (using similar triangles) or Hamiltonians to show that

$$\begin{aligned} \Delta\phi &= ga^2 \sin^4(\phi) \\ &= ga^2 \left( \underbrace{\frac{3}{8}}_{\text{Amplitude-dependent detuning}} - \underbrace{\frac{1}{2} \cos(2\phi) + \frac{1}{8} \cos(4\phi)}_{\text{Resonant driving}} \right) \end{aligned}$$

**Amplitude-dependent detuning:**  
doesn't depend on phase!

**Resonant driving:** periodic in  
betatron phase  $\phi$

- Useful (?) trick:

$$\sin^n(\phi) = \left( \frac{e^{i\phi} - e^{-i\phi}}{2i} \right)^n = \frac{1}{(2i)^n} \sum_{m=0}^n \binom{n}{m} (-1)^{(m+1)} (e^{i\phi})^{n-m} (e^{-i\phi})^m$$

binomial expansion

# Octupole Detuning Amplitude Dependence

- $\Delta\phi$  is an additional phase advance every turn
  - Dependent on amplitude  $a$  but not dependent on phase  $\phi$
- This is fundamentally a shift in the tune
  - Base (small-amplitude) tune is defined to be  $Q_0$
  - Tune of particles at amplitude  $a$  from octupoles is

$$Q = Q_0 + \frac{3}{16\pi}ga^2$$

- Nicely first order in octupole strength  $g$
- Turns out to be first order for quadrupoles, octupoles, dodecapoles, ... (you can see the pattern; dodecapole homework)
  - (Second order in nonlinearity strength for sextupoles, decapoles, ...)



## 10.2: Discrete Motion in $(J, \phi)$ Space

- Using action-angle space where  $J \equiv a^2/2$

$$Q = Q_0 + \frac{3}{8\pi} g J$$


- We can work out the general behavior in action along with phase to find general time evolution for **well-behaved particles**:

$$J_t = J_0 + \sum_{k=1}^{\infty} u_k \cos(k\phi_t + \phi_k) \quad (10.10)$$

$$\phi_t = \phi_0 + 2\pi Q t + \sum_{k=1}^{\infty} v_k \cos(k 2\pi Q t + \theta_k)$$

$u_k, v_k, \phi_k, \theta_k$  depend on nonlinearities  
 $J$  no longer constant

## 10.2: Discrete Motion in $(J, \phi)$ Space





- Recall Steve's comments on "smear", and Collins distortion functions (T.L. Collins, FNAL report 84/114)

# What is Really Happening Here?

- Tune varies with amplitude depending on nonlinearity

$$Q = Q_0 + \frac{3}{8\pi} gJ \quad \text{for octupoles}$$

- When  $Q_0$  is near resonance, particles with amplitude  $a_{\text{resonance}}$  have resonant tunes





# One-Turn Discrete “Hamiltonian”

- KAM surfaces suggest that we can write a “conserved” quantity and apply Hamiltonian dynamics to our discrete dynamical system

$$\Delta\phi = \frac{\partial H_1}{\partial J} \quad \Delta J = -\frac{\partial H_1}{\partial\phi}$$

- Here  $H_1$  is a “one-turn” discrete Hamiltonian. More generally we can include all nonlinearities:

$$H_1 = 2\pi(Q_{x0} J_x + Q_{y0} J_y) + \sum_{ijkl} V_{ijkl} J_x^{i/2} J_y^{j/2} \sin(k\phi_x + l\phi_y + \phi_{ijkl}) \quad (10.14)$$

Amplitude – dependent detuning when  $k = l = 0, i$  and/or  $j \neq 0$

# Motion Near Half-Integer Tunes

$$H_1 = 2\pi(Q_{x0} J_x + Q_{y0} J_y) + \sum_{ijkl} V_{ijkl} J_x^{i/2} J_y^{j/2} \sin(k\phi_x + l\phi_y + \phi_{ijkl}) \quad (10.14)$$

- One-turn maps from the one-turn “Hamiltonian” are still pretty jumpy
  - The fractional part of the tunes can be big even if everything else is perturbatively small
- But we can integrate the above equation and handwave an “N-turn” map
  - Near  $Q=k/N$  values, the phase advance is nearly  $2\pi$
  - All motion in N turns becomes perturbatively small

## Motion Near Half-Integer Tunes

$$H_1 = 2\pi(Q_{x0} J_x + Q_{y0} J_y) + \sum_{ijkl} V_{ijkl} J_x^{i/2} J_y^{j/2} \sin(k\phi_x + l\phi_y + \phi_{ijkl}) \quad (10.14)$$

$$Q = \frac{1}{2} + \delta Q \quad \delta Q \ll 1$$

$$H_2 = 2\pi \delta Q J + \left[ \frac{3}{8} - \frac{1}{2} \cos(2\phi) + \frac{1}{8} \cos(4\phi) \right] g J^2$$

or more generally, in the presence of many octupoles

$$H_2 = \underbrace{2\pi \delta Q J}_{\text{Tune difference from } 1/2} + \underbrace{[V_0 + V_2 \cos(2\phi + \phi_2) + V_4 \cos(4\phi + \phi_4)]}_{\text{octupole amplitude dependent detuning}} J^2$$

Tune difference  
from 1/2


octupole  
amplitude  
dependent  
detuning

Half-integer  
resonance  
driving


Quarter-integer  
resonance  
driving



# Motion Near Half-Integer Tunes: Figs 10.3-4



Normalised  
displacement,  $x$



Normalised horizontal angle,  $x'$

Can be used for slow extraction

# Entire USPAS courses on injection/extraction

- <http://uspas.fnal.gov/programs/2017/niu/courses/injection-extraction.shtml>

## Injection and Extraction of Beams

### Instructor:

Uli Wienands, Argonne National Lab and Edu Marin-Lacoma, CERN

---

### Purpose and Audience

This course provides an introduction to the physics and design of ring injection and extraction systems. Proton, ion, and electron systems will be covered. The course is appropriate for anyone with some background in accelerator physics and technology and with an interest in injection and extraction of beams, including operational staff.

- <http://uspas.fnal.gov/materials/17NIU/niu-injection-and-extraction.shtml>

## Course Materials - NIU - June 2017

### Injection and Extraction of Beams



---

course given by Uli Wienands, Argonne National Lab and Edu Marin-Lacoma, CERN

Updated pdf of the lecture hand-outs: [Accelerator Injection and Extraction](#)

Zipped archive of the Mad-X and Python scripts for the Mad-X injection-design exercise: [MADX Exercise files.zip](#) (Windows and Linux users can ignore the \_MACOSX folder that will be there after unzipping the file.)

# Half-Integer Slow Extraction





# Resonance Islands Revisited

- Todd's dissertation: E778 in the Fermilab Tevatron
- 5<sup>th</sup> order resonance islands driven to "second order" in sextupole strength
- Modern usage: resonance island extraction at CERN (Giovanozzi slides)

