

USPAS Graduate Accelerator Physics Homework 11

Due date: Wednesday February 10, 2021

1 Projection map Hamiltonian H_p

(10 points) Define the projection map P as linear motion R from a reference point to the location of a nonlinear magnet, where

$$R = \begin{pmatrix} c_x & s_x \\ -s_x & c_x \end{pmatrix} \quad c_x = \cos(\phi_x) \quad s_x = \sin(\phi_x)$$

followed by the nonlinear kick $\Delta x' = gx^n$, finally followed by inverse linear motion R^{-1} back to the reference point. Show that the discrete projection Hamiltonian representing P is given by

$$H_p = -\frac{g}{n+1}(c_x x + s_x x')^{n+1} \quad (1.1)$$

2 Hénon triangle near $Q = 1/3$

(10 points) Consider the equilateral triangle in (x, x') normalised phase space predicted by Equations 9.27 and 9.28.

- What is the radius of the largest circle that can be inscribed inside the triangle?
- What is the orientation of the triangle?
- What happens to the area and the orientation of the triangle as the tune Q is (slowly) swept through the value of $1/3$?

3 Hénon dynamic aperture simulation

(10 points) Use the simulation at <http://www.toddsatogata.net/2021-USPAS/lab/Henon.html> to investigate motion under the Hénon map by adjusting the two control parameters: tune Q and the number of turns tracked T . Launch multiple trajectories at many initial locations in phase space. Consider the plot of Hénon dynamic aperture (DA) versus tune shown in Figure 9.4.

- How small must $|Q - 1/3|$ be, for the triangular predictions of Exercise 9.4 (question 2 in this homework) to be reasonably valid?
- Devise and define a convenient quantitative measure of the size of the stable region – the DA. (There are many ways to do this.)
- How does the DA in the range $0.5 < Q < 1.0$ relate to the DA in the range below $Q = 0.5$? Why?